


Figure 1. Map of putative stocks, with individual assignments shown according to stock. Western Jones Sound (WJS), Baffin Bay (BB), Penny Strait-Lancaster Sound (PS-LIS), North Foxe Basin (N-FB), Central Foxe Basin (C-FB), Hudson Bay Davis Strait (HB-DS), Hudson Bay (HB), Devon Island (DI), Lancaster Sound (LS), the high Arctic cluster.



Walrus Harvesting Locations

Stewart, R.E.A. 2008. Redefining Walrus Stocks in Canada. *Arctic*. 61(3):292-308.




FIG. 5. Foxe Basin walrus harvesting locations for Igloolik and Hall Beach (Priest and Usher, 2004) and wintering areas (Born et al., 1995). Each kill symbol may represent more than one animal killed; N is the number of walrus reported by each community in the Priest and Usher (2004) database.





FIG. 6. Hudson Bay–Davis Strait walrus harvesting locations for Arviat, Cape Dorset, Chesterfield Inlet, Clyde River, Coral Harbour, Iqaluit, Kimmirut, Pangnirtung, Qikiqtarjuaq, Rankin Inlet, and Repulse Bay (Priest and Usher, 2004), Nunavik harvesting areas (interpreted from COSEWIC, 2006), and wintering areas (Born et al., 1995). Each kill symbol may represent more than one animal killed; N is the number of walrus reported by each community in the Priest and Usher (2004) database.

Foxe Basin

Figure 1: Foxe Basin walrus distribution



Coral Harbour (2014)

Figure 2: Coral Harbour walrus distribution






Figure 3: Coral Harbour walrus migration routes



Chesterfield Inlet (2008)

Figure 4: Chesterfield Inlet walrus distribution



Igloolik (2007)

Figure 5: Igloolik walrus distribution




Figure 6: Igloolik walrus distribution; observations from Interviewees 1-5





Figure 7: Igloolik walrus distribution; observations from Interviewees 6-11



Kimmirut (2010)

Figure 8: Kimmirut walrus distribution

